

# Chaos and Multifractals in the Solar Wind

**Wiesław M. Macek**

Faculty of Mathematics and Natural Sciences. College of Sciences,  
Cardinal Stefan Wyszyński University, Dewajtis 5, 01-815 Warsaw, Poland;  
Space Research Centre, Polish Academy of Sciences,  
Bartycka 18 A, 00-716 Warsaw, Poland  
e-mail: [macek@cbk.waw.pl](mailto:macek@cbk.waw.pl), <http://www.cbk.waw.pl/~macek>



# Abstract

The nature of the fluctuations in solar wind plasma parameters is still very little understood. The slow solar wind most likely originates from nonlinear processes in the solar corona. However, it appears that a certain kind of order does lie concealed within the irregular solar wind fluctuations, which can be described using methods of nonlinear time series analysis, based on fractal analysis and the theory of deterministic chaos. This involves the notions of fractal and multifractal sets, which could be presumably strange attractors in a certain state space of a given complex dynamical system. By employing the so-called false-nearest-neighbors method, we argue that the deterministic component of solar wind plasma dynamics should be low-dimensional. In fact, the results we have obtained using the method of topological embeddings indicate that the behavior of the solar wind can be approximately described by a low-dimensional chaotic attractor in the inertial manifold, which is a subspace of system phase space. We have also shown that the multifractal spectrum of the solar wind attractor is consistent with that for the multifractal measure of the self-similar generalized weighted Cantor set with two different scaling parameters describing nonuniform compression and one probability measure parameter of the chaotic attractor of the system. The values of the parameters fitted demonstrate small dissipation of the complex solar wind plasma and show that some parts of the attractor in phase space are visited much more frequently than other parts.

We hope that this generalized multifractal model will be a useful tool for analysis of intermittent turbulence in space plasmas. We thus believe that fractal analysis of chaotic systems could lead us to a deeper understanding of their nature, and maybe even to predict their seemingly unpredictable behavior.

**Within the complex dynamics of the solar wind's fluctuating intermittent plasma parameters, there is a detectable, hidden order described by a strange chaotic attractor that exhibits a multifractal structure.**


# Plan of Presentation

1. Introduction
  - Chaos and Fractals Basics
  - Importance of Chaos and Multifractality
2. Solar Wind Data
  - Solar Wind Fluctuations
  - Space Missions (Helios, Ulysses, ACE/WIND)
3. Methods for Solar Wind Attractor
  - Generalized Dimensions
4. Methods for Turbulence Models
  - Structure Functions Scaling (Energy Transfer Rate)
5. Results and Discussion
  - Attractor Reconstruction
  - Dimensions and Multifractality
  - Generalized Two-Scale Mutifractal Model for Intermittent Turbulence
  - Comparison with the usual p-model (Meneveau and Sreenivasan, 1987)
6. Conclusions

# Prologue

A **fractal** is a rough or fragmented geometrical object that can be subdivided in parts, each of which is (at least approximately) a reduced-size copy of the whole. Fractals are generally *self-similar* and independent of scale (fractal dimension).

A **multifractal** is a set of intertwined fractals. Self-similarity of multifractals is scale dependent (spectrum of dimensions).



Two-scale **Cantor** set.

# Chaos and Attractors

**CHAOS** ( $\chi\alpha\omicron\varsigma$ ) is

- APERIODIC long-term behavior
- in a DETERMINISTIC system
- that exhibits SENSITIVITY TO INITIAL CONDITIONS.

A positive finite Lyapunov exponent (metric entropy) implies chaos.

An **ATTRACTOR** is a *closed* set  $A$  with the properties:

1.  $A$  is an INVARIANT SET:  
any trajectory  $\mathbf{x}(t)$  that start in  $A$  stays in  $A$  for ALL time  $t$ .
2.  $A$  ATTRACTS AN OPEN SET OF INITIAL CONDITIONS:  
there is an open set  $U$  containing  $A$  ( $A \subset U$ ) such that if  $\mathbf{x}(0) \in U$ , then the distance from  $\mathbf{x}(t)$  to  $A$  tends to zero as  $t \rightarrow \infty$ .
3.  $A$  is MINIMAL:  
there is NO proper subset of  $A$  that satisfies conditions 1 and 2.

The nature of the fluctuations in solar wind plasma parameters is still very little understood. The slow solar wind most likely originates from nonlinear processes in the solar corona. However, it appears that a certain kind of order does lie concealed within the irregular solar wind fluctuations, which can be described using methods of nonlinear time series analysis, based on fractal analysis and the theory of deterministic chaos. This involves the notions of fractal and multifractal sets, which could be presumably strange attractors in a certain state space of a given complex dynamical system. By employing the so-called false-nearest-neighbors method, we argue that the deterministic component of solar wind plasma dynamics should be low-dimensional (Macek and Strumik, 2006). In fact, the results we have obtained using the method of topological embeddings indicate that the behavior of the solar wind can be approximately described by a low-dimensional chaotic attractor in the inertial manifold, which is a subspace of system phase space.

A direct determination of the solar wind attractor from the data is known to be a difficult problem. Indication for a chaotic attractor in the slow solar wind has been given by Macek (1998) and Macek and Redaelli (2000). In particular, Macek (1998) has calculated the correlation dimension of the reconstructed attractor in the solar wind and has provided tests for this measure of *complexity* including statistical surrogate data tests (Theiler *et al.*, 1992). Further, Macek and Redaelli (2000) have shown that the Kolmogorov entropy of the attractor is *positive* and finite, as it holds for a *chaotic* system.


The question of multifractality is of great importance because it allows us to look at intermittent turbulence in the solar wind (e.g., Marsch and Tu, 1997; Bruno *et al.*, 2001). Starting from Richardson's scenario of turbulence, many authors try to recover the observed scaling exponents, using some simple and more advanced fractal and multifractal models of turbulence describing distribution of the energy flux between cascading eddies at various scales. In particular, the multifractal spectrum has been investigated using Voyager (magnetic field) data in the outer heliosphere (e.g., Burlaga, 1991, 2001) and using Helios (plasma) data in the inner heliosphere (e.g., Marsch *et al.*, 1996). The multifractal scaling has also been tested using Ulysses observations (Horbury *et al.*, 1997) and with ACE/WIND data (e.g., Hnat *et al.*, 2003, 2007; Chapman *et al.*, 2006; Kiyani *et al.*, 2007).

We have also analyzed the spectrum for the solar wind attractor. This spectrum has been found to be consistent with that for the multifractal measure of the self-similar weighted baker's map with two parameters describing uniform compression and natural invariant probability measure of the attractor of the system (Macek, 2002, 2003, 2006, 2007; Macek et al., 2005, 2006).


Therefore, here in order to further quantify the multifractality, we also consider the generalized weighted Cantor set with two different scales describing nonuniform compression in the context of turbulence cascade. Even though one can find this general Cantor set in many classical textbooks, e.g., (Falconer, 1952, Ott, 1993), it is still difficult to understand this strange attractor that exhibits multifractality in various complex real systems, also in case of intermittent turbulence.

Hence we argue that there is, in fact, need to use a two-scale cascade model. Therefore, we investigate the resulting multifractal spectrum of the energy flux depending on two scaling parameters and one probability measure parameter, demonstrating that intermittent pulses are stronger for asymmetric scaling and a better agreement is obtained especially for  $q < 0$  (Macek, 2007). We hope that this generalized new asymmetric multifractal model could shed light on the nature of turbulence and will be a useful tool for analysis of intermittent turbulence in various environments.

- W. M. Macek, *Physica D* **122**, 254–264 (1998).
- W. M. Macek and S. Redaelli, *Phys. Rev. E* **62**, 6496–6504 (2000).
- W. M. Macek, Multifractality and chaos in the solar wind, in *Experimental Chaos*, edited by S. Boccaletti, B. J. Gluckman, J. Kurths, L. M. Pecora, and M. L. Spano, American Institute of Physics, New York, 2002, Vol. 622, pp. 74–79.
- W. M. Macek, The multifractal spectrum for the solar wind flow, in *Solar Wind 10*, edited by M. Velli, R. Bruno, F. Malara, American Institute of Physics, New York, 2003, vol. 679, pp. 530–533.
- W. M. Macek, R. Bruno, G. Consolini, Generalized dimensions for fluctuations in the solar wind, *Phys. Rev. E* **72**, 017202 (2005).
- W. M. Macek, R. Bruno, G. Consolini, Testing for multifractality of the slow solar wind, *Adv. Space Res.* **37**, 461-466 (2006).
- W. M. Macek, Modeling multifractality of the solar wind, *Space Sci. Rev.* **122**, 329–337 (2006).
- W. M. Macek, Multifractality and intermittency in the solar wind, *Nonlinear Processes in Geophysics*, 14, 1–7 (2007).
- W. M. Macek, A. Szczepaniak, Generalized Two-Scale Weighted Cantor Set Model for Solar Wind Turbulence submitted (2007).



A schematic model of the solar wind "ballerina": the Sun's two hemispheres are separated by a neutral layer of a form reminiscent of a 'ballerina's skirt'. In the inner heliosphere the solar wind streams are of two forms called the slow ( $\approx 400 \text{ km s}^{-1}$ ) and fast ( $\approx 700 \text{ km s}^{-1}$ ). The fast wind is associated with coronal holes and is relatively uniform and stable, while the slow wind is quite variable, taken from (Schwenn and Rosenbauer, 1984).




Structures in the solar wind and their sources in the corona (solar map), taken from (Schwenn and Rosenbauer, 1984).

# Helios Spacecraft



Orbits



# Methods of Data Analysis

## Generalized Dimensions

The generalized dimensions are important characteristics of *complex* dynamical systems. Since these dimensions are related to frequencies with which typical orbits in phase space visit different regions of the system, they can provide information about its dynamics.

The modern technique of nonlinear time series analysis allows to estimate the multifractal measure directly from a single time series.

## Structure Functions Scaling

$S_u^q(l)$ ,  $q$ th order structure function ( $q > 0$ ) in the inertial range ( $\eta \ll l \ll L$ )

$$S_u^q(l) = \langle |u(x+l) - u(l)|^q \rangle \sim l^{\xi(q)} \quad (1)$$

$u(x)$ , a velocity component parallel to the  $l$

$\xi(q)$ , a scaling exponent.

## Energy Transfer Rate

$$\varepsilon_l \sim \frac{S_u^3(l)}{l} \quad \mu_i = \frac{\varepsilon_l}{\langle \varepsilon_L \rangle} \quad (2)$$

$$\sum_i \mu_i^q \sim l^{\tau(q)} \quad (3)$$

$$\tau(q) = (q-1) D_q \quad (4)$$


$\varepsilon_l$ , energy transfer rate

$\mu_i$ , probability measure of  $i$ th eddy in the  $d$ -dimensional physical space.

From Equations (1 ) to (4) we have (Tsang *et al.*, 2005):

$$D_q = d + \frac{\xi(3q) - q\xi(3)}{q-1} \quad (5)$$

# Mutifractal Models for Turbulence



**Fig. 1.** Generalized two-scale Cantor set model for turbulence (Macek, 2007).

For the generalized self-similar weighted Cantor set (acting on the unit interval) we use the following partition function at  $n$ -th level of construction (Hentschel and Procaccia 1983; Halsey *et al.*, 1986)

$$\Gamma_n(l_1, l_2, p_1, p_2) = \left( \frac{p_1^q}{l_1^{\tau(q)}} + \frac{p_2^q}{l_2^{\tau(q)}} \right)^n = 1 \quad (6)$$

Parameters:

- $p_1 = p \leq 1/2$ , natural invariant measure on the attractor of the system, the probability of visiting one region of the interval (the probability of visiting the remaining region is  $p_2 = 1 - p$ );
- $l_1 + l_2 \leq 1$ , two nonuniform compression (dissipation) parameters (stretching and folding in the phase space).

# Solutions

**Transcendental equation** (for  $n \rightarrow \infty$ )

$$p^q l_1^{-\tau(q)} + (1-p)^q l_2^{-\tau(q)} = 1 \quad (7)$$

**Legendre transformation**

$$\alpha(q) = \frac{d \tau(q)}{dq} \quad (8)$$

$$f(\alpha) = q\alpha(q) - \tau(q) \quad (9)$$

For  $l_1 = l_2 = s$  and any  $q$  in Eq. (7) one has for the generalized dimension of the attractor (projected onto one axis)

$$(q-1)D_q = \frac{\ln[p^q + (1-p)^q]}{\ln s}. \quad (10)$$

No dissipation ( $s = 1/2$ ):  
the multifractal cascade  $p$ -model for fully developed turbulence,  
the generalized weighted Cantor set (Meneveau and Sreenivasan, 1987).  
The usual middle one-third Cantor set (without any multifractality):  
 $p = 1/2$  and  $s = 1/3$ .

The difference of the maximum and minimum dimension  
(the least dense and most dense points on the attractor)

$$D_{-\infty} - D_{+\infty} = \frac{\ln(1/p - 1)}{\ln(1/s)} \quad (11)$$


In the limit  $p \rightarrow 0$  this difference rises to infinity (degree of multifractality).

# Data



Table 1: The time intervals of Helios 2 data in 1976 for slow and fast solar wind streams measured at various distances from the Sun.

	~ 0.3 AU	~ 0.97 AU
Slow streams	099:00:00:29 - 099:23:34:27	026:00:00:33 - 026:23:58:45
	100:00:09:57 - 100:23:59:41	027:00:00:05 - 027:23:59:57
ddd:hh:mm:ss	101:00:00:21 - 101:23:59:47	028:00:00:39 - 028:23:59:59
	102:00:00:27 - 102:23:59:31	029:00:00:41 - 029:23:59:23
Fast streams	105:00:00:31 - 105:23:59:25	021:00:00:09 - 021:23:59:41
	106:00:00:45 - 106:23:59:35	022:00:00:17 - 022:23:59:51
ddd:hh:mm:ss	107:00:00:17 - 107:23:59:49	023:00:00:31 - 023:23:59:35
	108:00:00:29 - 108:23:59:55	024:00:00:07 - 024:23:59:43



# Attractor Reconstruction




**Fig. 2.** The projection of the attractor onto the three-dimensional space, reconstructed from the detrended data,  $T = 4 \Delta t$ , using (a) the moving average and also (b) the singular-value decomposition filters ( $\Psi = U$ ), taken from (Macek, 1998).




**Fig. 3.** (a) The generalized dimensions  $D_q$  in Equation (4) as a function of  $q$ . The correlation dimension is  $D_2 = 3.4 \pm 0.1$ . The values of  $D_q + 3$  are calculated analytically for one-scale weighted Cantor set (baker's map) with  $p = 0.12$  and  $s = 0.47$  (dashed line). (b) The singularity spectrum  $f(\alpha)$  as a function of  $\alpha$ . The values of  $f(\alpha)$  projected onto one axis for the weighted baker's map with the same parameters (dashed line), taken from (Macek, 2006).




**Fig. 4.** (a) The generalized dimensions  $D_q$  in Equation (4) as a function of  $q$ . The values of  $D_q + 3$  are calculated analytically for the weighted two-scale Cantor set with  $p = 0.20$  and  $l_1 = 0.60$ ,  $l_2 = 0.25$  (dashed line). (b) The singularity spectrum  $f(\alpha)$  as a function of  $\alpha$ . The values of  $f(\alpha)$  projected onto one axis for the weighted two-scale Cantor set with the same parameters (dashed line), taken from (Macek, 2007).




**Fig. 5.** The generalized dimensions  $D_q$  as a function of  $q$ . The values for one-dimensional turbulence are calculated for the generalized two-scale (continuous lines) model and the usual one-scale (dashed lines)  $p$ -model and fitted using the  $u = v_x$  radial velocity components (diamonds) for the slow (a) and fast (b) solar wind streams at distances of 0.3 AU (Macek and Szczepaniak, 2007).




**Fig. 5.** The generalized dimensions  $D_q$  as a function of  $q$ . The values for one-dimensional turbulence are calculated for the generalized two-scale (continuous lines) model and the usual one-scale (dashed lines)  $p$ -model and fitted using the  $u = v_x$  radial velocity components (diamonds) for the slow (c) and fast (d) solar wind streams at distances of 0.97 AU (Macek and Szczepaniak, 2007).



**Fig. 6.** The singularity spectrum  $f(\alpha)$  as a function of  $\alpha$ . The values for one-dimensional turbulence are calculated for the generalized two-scale (continuous lines) model and the usual one-scale (dashed lines)  $p$ -model and fitted using the  $u = v_x$  radial velocity components (diamonds) for the slow (a) and fast (b) solar wind streams at distances of 0.3 AU (Macek and Szczepaniak, 2007).



**Fig. 6.** The singularity spectrum  $f(\alpha)$  as a function of  $\alpha$ . The values for one-dimensional turbulence are calculated for the generalized two-scale (continuous lines) model and the usual one-scale (dashed lines)  $p$ -model and fitted using the  $u = v_x$  radial velocity components (diamonds) for the slow (c) and fast (d) solar wind streams at distances of 0.97 AU (Macek and Szczepaniak, 2007).



**Fig. 7.** The multifractal measure  $\epsilon/\langle\epsilon\rangle$  on the unit interval for (a) the usual one-scale  $p$ -model (Meneveau and Sreenivasan, 1987) and (b) the generalized two-scale cascade model. Intermittent pulses are stronger for the model with two different scaling parameters (Macek and Szczepaniak, 2007).

The value of parameter  $p$  (within some factor) is related to the usual models, which starting from Richardson's scenario of turbulence, try to recover the observed scaling exponents, which is based on the  $p$ -model of turbulence (e.g. Meneveau and Sreenivasan, 1987).

The value of  $p \approx 0.2$  obtained here is roughly consistent with the fitted value in the literature both for laboratory and the solar wind turbulence, which is in the range  $0.13 \leq p \leq 0.3$  (e.g., Burlaga, 1991; Carbone, 1993; Carbone and Bruno, 1996; Marsch *et al.*, 1996).

One should only bear in mind that here we take probability measure directly on the solar wind attractor, which quantifies multifractal nonuniformity of visiting various parts of the attractor in the phase space, while the usual  $p$ -model is related to the solar wind turbulence cascade for the energy transfer rate, which resides in the physical space.

# Conclusions

In this way, we have supported our conjecture that

- trajectories describing the system in the inertial manifold of phase space asymptotically approach the attractor of low-dimension (Macek, 1998).
- The obtained multifractal spectrum of this attractor is consistent with that for the multifractal measure on the generalized weighted two-scale Cantor set, which is a strange attractor that exhibits stretching and folding properties leading to sensitive dependence on initial conditions (Macek, 2006, 2007).
- The values of the parameters fitted demonstrates small dissipation of the complex solar wind dynamical system and shows that some parts of the attractor in phase space are visited at least one order of magnitudes more frequently than other parts.

- We have also studied departure from Kolmogorov scaling indicating multifractal (intermittent) behavior of the solar wind in the inner heliosphere.
  - We confirm that the degree of multifractality of the solar wind is different for slow and fast streams.
  - Our analysis shows that slow solar wind velocity fluctuations are more intermittent and more anisotropic than for the fast solar wind.
  - Also as the heliocentric distance increases the solar wind becomes more multifractal consistent with other studies. In particular, we observe radial evolution of multifractality (intermittency) as noticed, e.g., by Marsch *et al.* (1996).
- Basically, the generalized dimensions for solar wind are consistent with the generalized  $p$ -model for both positive and negative  $q$ , but rather with different scaling parameters for sizes of eddies, while the usual  $p$ -model can only reproduce the spectrum for  $q \geq 0$ .  
 We show that intermittent pulses are stronger for the model with asymmetric scaling and a somewhat better agreement with the solar wind data is obtained.
- Therefore, we propose the generalized two-scale weighted Cantor set model describing intermittent energy transfer cascade for analysis of space plasma turbulence.

One can expect that the attractor in the low-speed solar wind plasma should contain information about the dynamic variations of the coronal streamers. It is also possible that it represents a structure of the time sequence of near-Sun coronal fine-stream tubes.

# Epilogue

Thus these results provide supporting evidence for **multifractal** structure of the solar wind in the inner heliosphere.

This means that the observed **intermittent** behavior of the solar wind's velocity and Alfvénic fluctuations results from intrinsic *nonlinear* dynamics rather than from random external forces.

The multifractal structures, convected by the solar wind, might probably be related to the complex topology shown by the magnetic field at the source regions of the solar wind.

Thank you!



# References

- [Bruno et al.(2001)] Bruno, R., V. Carbone, P. Veltri, E. Pietropaolo, and B. Bavassano (2001), Identifying intermittent events in the solar wind, *Planet. Space Sci.*, *49*, 1201–1210.
- [Burlaga(1991)] Burlaga, L. F. (1991), Multifractal structure of the interplanetary magnetic field: Voyager 2 observations near 25 AU, 1987–1988, *Geophys. Res. Lett.*, *18*, 69–72.
- [Burlaga(2001)] Burlaga, L. F. (2001), Lognormal and multifractal distributions of the heliospheric magnetic field, *J. Geophys. Res.*, *106*, 15917–15927.
- [Carbone(1993)] Carbone, V. (1993), Cascade model for intermittency in fully developed magnetohydrodynamic turbulence, *Phys. Rev. Lett.*, *71*, 1546–1548.
- [Carbone and Bruno(1996)] Carbone, V., and R. Bruno (1996), Cancellation exponents and multifractal scaling laws in the solar wind magnetohydrodynamic turbulence, *Ann. Geophys.*, *14*, 777–785.
- [Halsey et al.(1986)] Halsey, T. C., M. H. Jensen, L. P. Kadanoff, I. Procaccia, and B. I. Shraiman (1986), Fractal measures and their singularities: The characterization of strange sets, *Phys. Rev. A*, *33*, 1141–1151.
- [Hentschel and Procaccia(1983)] Hentschel, H. G. E., and I. Procaccia (1983), The infinite number of generalized dimensions of fractals and strange attractors, *Physica D*, *8*, 435–444.
- [Macek(1998)] Macek, W. M. (1998), Testing for an attractor in the solar wind flow, *Physica D*, *122*, 254–264.
- [Macek(2002)] Macek, W. M. (2002), Multifractality and chaos in the solar wind, edited by S. Boccaletti, B.J. Gluckman, J. Kurths, L.M. Pecora, M.L. Spano, *Experimental Chaos*, *622*. American Institute of Physics, New York, pp. 74–79.
- [Macek(2003)] Macek, W. M. (2003), The multifractal spectrum for the solar wind flow, edited by R. Velli, R. Bruno, and F. Malara, *Solar Wind 10*, *679*. American Institute of Physics, New York, pp. 530–533.

- [Macek(2006)] Macek, W. M. (2006), Modeling multifractality of the solar wind, *Space Sci. Rev.*, 122, 329–337, doi:10.1007/s11214-006-8185-z.
- [Macek(2007)] Macek, W. M. (2007), Multifractality and intermittency in the solar wind, *Nonlinear Proc. Geophys.* 14, 1–7.
- [Macek and Redaelli(2000)] Macek W. M., and S. Redaelli (2000), Estimation of the entropy of the solar wind flow, *Phys. Rev. E*, 62, 6496–6504.
- [Macek and Szczepaniak (2007)] Macek, W. M., and A. Szczepaniak (2007), Asymmetric Multifractal Model for Intermittent Turbulence, submitted.
- [Macek et al.(2005)] Macek W. M., R. Bruno, and G. Consolini (2005), Generalized dimensions for fluctuations in the solar wind, *Phys. Review E*, 72, 017202, doi: 10.1103/PhysRevE.72.017202;
- [Macek et al.(2006)] Macek W. M., R. Bruno, and G. Consolini (2006), Testing for multifractality of the slow solar wind, *Adv. Space Res.*, 37, 461–466, doi:10.1016/j.asr.2005.06.057.
- [Mandelbrot(1989)] Mandelbrot, B. B. (1989), Multifractal measures, especially for the geophysicist, in *Pure and Applied Geophys.*, 131, Birkhäuser Verlag, Basel, pp. 5–42.
- [Meneveau and Sreenivasan(1987)] Meneveau, C., and K. R. Sreenivasan (1987), Simple multifractal cascade model for fully developed turbulence, *Phys. Rev. Lett.*, 59, 1424–1427.
- [Marsch and Tu(1997)] Marsch, E., and C.-Y Tu (1997), Intermittency, non-Gaussian statistics and fractal scaling of MHD fluctuations in the solar wind, *Nonlinear Proc. Geophys.*, 4, 101–124.
- [Marsch et al.(1996)] Marsch, E., C.-Y. Tu, and H. Rosenbauer (1996), Multifractal scaling of the kinetic energy flux in solar wind turbulence, *Ann. Geophys.*, 14, 259–269.
- [Ott(1993)] Ott, E. (1993), *Chaos in Dynamical Systems*, Cambridge University Press, Cambridge.
- [Schwenn(1990)] Schwenn, R. (1990), Large-scale structure of the interplanetary medium, edited by Schwenn, R., and E. Marsch, *Physics of the Inner Heliosphere*, 20. Springer-Verlag, Berlin, pp. 99–182.

[Tsang et al. (2005)] Tsang, Yue-Kin, E. Ott, T. M. Antonsen, Jr., and P. N. Guzdar (2005), Intermittency in two-dimensional turbulence with drag, *Phys. Rev. E*, 71, 066313, doi: 10.1103/PhysRevE.71.066313.